

Relativistic diffusion model and analysis of large transverse momentum distributions

Naomichi Suzuki* and Minoru Biyajima†

*Department of Comprehensive Management, Matsumoto University, Matsumoto 390-1295, Japan

†Department of Physics, Shinshu University, Matsumoto, 390-8621, Japan

Abstract. In order to describe large transverse momentum (p_T) distributions observed in high energy nucleus-nucleus collisions, a stochastic model in the three dimensional rapidity space is introduced. The fundamental solution of the radial symmetric diffusion equation is Gaussian-like in radial rapidity. We can also derive a p_T or radial rapidity distribution function, where a distribution of emission center is taken into account. It is applied to the analysis of observed large p_T distributions of charged particles. It is shown that our model approaches to a power function of p_T in the high transverse momentum limit.

Keywords: Relativistic heavy ion collisions, Stochastic model, Large p_T distributions

PACS: 25.75.-q, 02.50.Ey

INTRODUCTION

In high energy nucleus-nucleus collisions, colliding energy $\sqrt{s_{NN}}$ per nucleon grows up to $\sqrt{s_{NN}} = 200$ GeV, and secondary particles with transverse momentum p_T more than 10 GeV/c are observed. The observed p_T distributions have long tail compared with exponential distribution in p_T .

As is well known, the fundamental solution of stochastic process is Gaussian if variables in the Euclidian space are used. Therefore, as long as we consider stochastic equations in the transverse momentum space, it is very hard to describe observed p_T distributions. This fact suggests that a relativistic approach to the stochastic process would be needed.

In Ref. [1], an empirical formula for large p_T distributions at polar angle $\theta = \pi/2$,

$$E \frac{d^3\sigma}{d^3p} \Big|_{\theta=\pi/2} = A \exp[-y_T^2/(2L_T)],$$

$$y_T = \frac{1}{2} \ln \frac{E + |\mathbf{p}_T|}{E - |\mathbf{p}_T|}, \quad (1)$$

was proposed from the analogy of Landau's hydrodynamical model. Polar angle θ ($0 \leq \theta \leq \pi$) is measured from the beam direction of colliding nuclei or incident particles. In Eq. (1), E denotes energy of an observed particle, L_T is a parameter, and y_T is called the "transverse rapidity". Equation (1) well describes large p_T distributions for $p + p \rightarrow \pi^0 + X$ and $p + p \rightarrow \pi^\pm + X$. However, it cannot be derived from the hydrodynamical model.

As for the relativistic approach to the stochastic equation, we consider the diffusion equation in the three dimensional rapidity space or Lobachevsky space, which is non-

Euclidean. In order to specify longitudinal and transverse expansions, it would be appropriate to consider the diffusion equation in the geodesic cylindrical coordinate, where longitudinal rapidity y , transverse rapidity ξ and azimuthal angle ϕ are used. The longitudinal rapidity y , and the transverse rapidity ξ are defined respectively as,

$$y = \ln \frac{E + p_L}{m_T}, \quad \xi = \ln \frac{m_T + |\mathbf{p}_T|}{m},$$

where E , p_L , $\mathbf{p}_T$ and m denote energy, longitudinal momentum, transverse momentum, and mass of the observed particle, respectively, and $m_T = \sqrt{\mathbf{p}_T^2 + m^2}$.

The diffusion equation in the geodesic cylindrical coordinate is given by,

$$\frac{\partial f}{\partial t} = \frac{D}{\cosh^2 \xi} \frac{\partial^2 f}{\partial y^2} + \frac{D}{\sinh \xi \cosh \xi} \frac{\partial}{\partial \xi} \left(\sinh \xi \cosh \xi \frac{\partial f}{\partial \xi} \right) + \frac{D}{\sinh^2 \xi} \frac{\partial^2 f}{\partial \phi^2}, \quad (2)$$

where D denotes a diffusion constant. However, we cannot solve Eq.(2) analytically at present. Therefore, we should consider somewhat simpler case.

We have proposed the relativistic diffusion model, and analyzed large p_T distributions for charged particles in $Au + Au$ collisions [2], where a radial flow effect is not included.

In section 2, the relativistic diffusion model is briefly explained. A distribution of the initial radial rapidity is taken into account. It would correspond to the distribution of radial flow. In section 3, large p_T distributions for charged particles observed at RHIC [3, 4] are analyzed. The magnitude of radial flow is estimated from p_T distributions. In section 4, high transverse momentum limit of our model is taken and relation to Hagedorn's model for p_T distribution inspired by the QCD is discussed. Final section is devoted to summary and discussions.

DIFFUSION EQUATION WITH RADIAL SYMMETRY IN THE THREE DIMENSIONAL RAPIDITY SPACE

For simplicity, we consider the diffusion equation with radial symmetry in the geodesic polar coordinate system,

$$\frac{\partial f}{\partial t} = \frac{D}{\sinh^2 \rho} \frac{\partial}{\partial \rho} \left(\sinh^2 \rho \frac{\partial f}{\partial \rho} \right), \quad (3)$$

with an initial condition,

$$f(\rho, t = 0) = \frac{\delta(\rho - \rho_0)}{4\pi \sinh^2 \rho}. \quad (4)$$

In Eq. (3), ρ denotes the radial rapidity, which is written with energy E , momentum $\mathbf{p}$ and mass m of observed particle as,

$$\rho = \ln \frac{E + |\mathbf{p}|}{m}. \quad (5)$$

Inversely, energy and momentum are written respectively as

$$E = m \cosh \rho, \quad |\mathbf{p}| = \sqrt{p_L^2 + \mathbf{p}_T^2} = m \sinh \rho. \quad (6)$$

The solution of Eq. (3) with the initial condition (4) is given [5] by

$$f(\rho, \rho_0, t) = \frac{1}{2\pi\sqrt{4\pi Dt}} e^{-Dt} \frac{\sinh \frac{\rho_0 \rho}{2\pi Dt}}{\sinh \rho_0 \sinh \rho} \exp\left[-\frac{\rho^2 + \rho_0^2}{4Dt}\right]. \quad (7)$$

From Eq.(7), the following equation is obtained;

$$f(\rho, t) = \lim_{\rho_0 \rightarrow 0} f(\rho, \rho_0, t) = (4\pi Dt)^{-3/2} e^{-Dt} \frac{\rho}{\sinh \rho} \exp\left[-\frac{\rho^2}{4Dt}\right]. \quad (8)$$

In Eq. (7), ρ_0 denotes the radial rapidity of an emission center. It would be identified to the radial flow rapidity. We have estimated radial flow rapidity using Eq. (7) [6].

If the distribution of ρ_0 is taken into account, it would be reasonable to assume that it distributes randomly with dispersion σ_0^2 ;

$$f(\rho_0, t) = (2\pi\sigma_0^2)^{-3/2} e^{-\sigma_0^2/2} \frac{\rho_0}{\sinh \rho_0} \exp\left[-\frac{\rho_0^2}{2\sigma_0^2}\right].$$

Then the distribution function of radial rapidity is given by the following equation,

$$\begin{aligned} f_c(\rho, t) &= \int_0^\infty f(\rho, \rho_0, t) f(\rho_0, t) \sinh^2 \rho_0 \sin \theta d\rho_0 d\theta d\phi \\ &= (2\pi\sigma_T^2)^{-3/2} e^{-\sigma_T^2/2} \frac{\rho}{\sinh \rho} \exp\left[-\frac{\rho^2}{2\sigma_T^2}\right], \end{aligned} \quad (9)$$

$$\sigma_T^2 = 2Dt + \sigma_0^2. \quad (10)$$

Equation (9) is obtained from Eq. (8), if $2Dt$ is replaced by σ_T^2 . Parameter σ_T^2 is connected to the moment of ρ_0 as,

$$\langle \rho_0^2 \rangle = \int_0^\infty \rho_0^2 f(\rho_0, t) \sinh^2 \rho_0 \sin \theta d\rho_0 d\theta d\phi = \sigma_0^4 + 3\sigma_0^2.$$

When $\rho \ll 1$, $|\mathbf{p}| = m \sinh \rho \simeq m\rho$. Then, Eq. (8) reduces to

$$f(\rho, t) \simeq \exp[-\rho^2/(2\sigma(t)^2)], \quad \sigma(t)^2 = 2Dt. \quad (11)$$

If we assume that Eq. (11) should coincide with the Maxwell-Boltzmann distribution, we have an identity, $k_B T = m\sigma(t)^2$, where k_B is the Boltzmann constant. Then, Eq. (10) reduces to

$$\sigma_T^2 = k_B T / m + \left(-\frac{3}{2} + \frac{1}{2} \sqrt{9 + 4\langle \rho_0^2 \rangle}\right). \quad (12)$$

When, $\sigma_0^2 \ll 1$ or $\rho_0^2 \ll 1$, Eq. (10) is written as,

$$\sigma_T^2 \simeq k_B T / m + \langle \rho_0^2 \rangle / 3. \quad (13)$$

ANALYSIS OF LARGE P_T DISTRIBUTIONS OF CHARGED PARTICLES

Transverse momentum distributions of charged particles observed by STAR [3] and PHENIX [4] Collaborations are analyzed by Eq. (9). Data are taken at $\theta = \pi/2$. In this case, the identity, $\rho = y_T = \xi$, is satisfied.

The results on the data by STAR Collaboration are shown in Fig. 1 and Table 1.




FIGURE 1. (a) p_T distribution for $Au + Au \rightarrow (h^+ + h^-)/2 + X$ at $y = 0$ [3], and (b) centrality dependence of $\langle \rho_0^2 \rangle^{1/2}$ extracted from p_T distributions

TABLE 1. Parameters on p_T distributions estimated by Eq. (9) in $Au + Au \rightarrow (h^+ + h^-)/2 + X$ at $y = 0$ at $\sqrt{s_{NN}} = 200$ GeV [3]

centrality	C	σ_T^2	m	$\chi^2_{\min}/\text{n.d.f}$
00-05%	1126.3 ± 51.0	0.332 ± 0.003	0.570 ± 0.007	244.9/32
05-10%	916.2 ± 47.5	0.332 ± 0.004	0.567 ± 0.009	128.8/32
10-20%	780.0 ± 36.2	0.351 ± 0.004	0.536 ± 0.008	144.2/32
20-30%	590.1 ± 28.9	0.360 ± 0.004	0.519 ± 0.008	98.1/32
30-40%	451.5 ± 22.7	0.374 ± 0.004	0.492 ± 0.008	95.4/32
40-60%	300.99 ± 14.47	0.402 ± 0.004	0.442 ± 0.006	51.7/32
60-80%	150.91 ± 7.43	0.444 ± 0.004	0.369 ± 0.006	26.5/32
pp(NSD)	17.654 ± 1.424	0.455 ± 0.006	0.324 ± 0.008	16.1/29

Information on radial flow is extracted by the use of Eq. (12) under the assumption that the hadronization temperature is constant irrespective of centrality. The results are shown in Fig.1b. The mean radial flow rapidity, $\langle \rho_0^2 \rangle^{1/2}$, depends on centrality very weakly, if it is less than 60%. At $k_B T = 0.13$ GeV, $\langle \rho_0^2 \rangle^{1/2} \simeq 0.57$ for centrality $< 60\%$.

The results on the data by FHENIX Collaboration are shown in Fig. 2 and Table 2. Estimated value of $\langle \rho_0^2 \rangle^{1/2}$ is somewhat larger than that from STAR Collaboration at the same temperature.




FIGURE 2. (a) p_T distributions for $Au + Au \rightarrow (h^+ + h^-)/2 + X$ at $y = 0$ [4], and (b) centrality dependence of $\langle \rho_0^2 \rangle^{1/2}$ extracted from p_T distributions

TABLE 2. Parameters on p_T distributions estimated by Eq. (9) in $Au + Au \rightarrow (h^+ + h^-)/2 + X$ at $y = 0$ at $\sqrt{s_{NN}} = 200$ GeV [4]

centrality	C	σ_T^2	m	$\chi^2_{min}/n.d.f$
00-05%	410.00 ± 42.99	0.277 ± 0.007	0.745 ± 0.023	32.1/31
00-10%	408.50 ± 41.25	0.284 ± 0.007	0.726 ± 0.021	35.5/31
10-20%	305.21 ± 30.84	0.293 ± 0.007	0.708 ± 0.021	26.4/31
20-30%	265.03 ± 27.18	0.313 ± 0.007	0.653 ± 0.019	27.8/31
30-40%	209.37 ± 21.82	0.330 ± 0.007	0.611 ± 0.019	16.8/29
40-50%	143.61 ± 15.42	0.338 ± 0.007	0.587 ± 0.018	15.4/29
50-60%	103.03 ± 11.65	0.358 ± 0.008	0.537 ± 0.018	9.9/29
60-70%	70.343 ± 8.419	0.380 ± 0.009	0.485 ± 0.017	5.1/29
70-80%	40.688 ± 5.284	0.390 ± 0.010	0.457 ± 0.018	7.4/28
80-92%	22.286 ± 3.126	0.406 ± 0.011	0.419 ± 0.018	10.3/27

HIGH TRANSVERSE MOMENTUM LIMIT OF THE MODEL

In the 1970's, when large p_T distributions are observed in accelerator experiments, many models, which have power-law behavior in p_T are proposed. In Ref. [7], a model for p_T distribution, inspired by the QCD, is proposed as $(p_0/(p_T + p_0))^n$. It approaches an exponential distribution of p_T for $p_T \rightarrow 0$, and a power function of p_T for $p_T \rightarrow \infty$.

In this section, high transverse momentum limit of Eq. (9) is examined at $\theta = \pi/2$, where the identity $\rho = \ln((m_T + p_T)/m)$ holds. Radial rapidity contained in the Gaussian-like part in Eq. (9) is rewritten as,

$$\exp\left[-\frac{\rho^2}{2\sigma_T^2}\right] = \left(\frac{m_T + p_T}{m}\right)^{-\rho/(2\sigma_T^2)}. \quad (14)$$

For any positive number ε , we have,

$$\lim_{p_T \rightarrow \infty} \frac{\rho}{p_T^\varepsilon} = \lim_{p_T \rightarrow \infty} \frac{1}{p_T^\varepsilon} \ln \left(\frac{m_T + p_T}{m} \right) = 0.$$

Therefore, we can approximate ρ in the exponent in Eq. (14) as a constant, which is written by $2c_0$, within some finite transverse momentum range. Then Eq. (9) reduces to,

$$f_c(\rho, t) \sim \left(\frac{m}{m_T + p_T} \right)^{c_0/\sigma_T^2} \frac{m}{p_T} \sim \left(\frac{m}{2p_T} \right)^{c_0/\sigma_T^2+1}. \quad (15)$$

From Eq. (15), one can see that Eq. (9) shows the power-law behavior in the high transverse momentum limit, and that the power becomes smaller as σ_T^2 , which should increase with the colliding energy $\sqrt{s_{NN}}$, increases.

SUMMARY AND DISCUSSIONS

In order to analyze large p_T distributions of charged particles observed at RHIC, the relativistic stochastic process in the three dimensional rapidity space, which is non-Euclidean, is introduced. The solution is Gaussian-like in radial rapidity, where the radial flow rapidity ρ_0 is included. It is very similar to the formula proposed in Ref. [1] at $\theta = \pi/2$. (See also Ref. [8].)

Transverse momentum distributions for charged particles in $Au + Au$ collisions at $y = 0$ at $\sqrt{s_{NN}} = 200$ GeV are analyzed. From the observed p_T distributions, an effect of radial flow is subtracted by the use of assumption that the hadronization temperature does not depend on the centrality.

The averaged radial flow rapidity, $\langle \rho_0^2 \rangle^{1/2}$, depends on the centrality very weakly from 0% up to 60%. It decreases rapidly, as the centrality increases above 60%.

We have derived that our formula (9) shows power-law behavior in p_T in the high transverse momentum limit. Therefore, it behaves like the Gaussian distribution in p_T when $p_T << m$, and like the power-law distribution in p_T when $p_T \gg 1$.

ACKNOWLEDGMENTS

Authors would like to thank RCNP at Osaka University, Faculty of Science, Shinshu University, and Matsumoto University for financial support.

REFERENCES

1. M. Duong-van and P. Carruthers, *Phys. Rev. Letters*, **31**, 133 (1973)
2. N. Suzuki and M. Biyajima, *Acta Phys. Polon.* **B35**, 283 (2004); hep-ph/0404112
3. J. Adams, et al., STAR Collaboration, *Phys. Rev. Letters*, **91**, 172301 (2003)
4. S.S. Adler, et al., PHENIX Collaboration, *Phys. Rev.*, **C69**, 034910 (2004)
5. N. Suzuki and M. Biyajima, math-ph/0406040, unpublished
6. N. Suzuki and M. Biyajima, hep-ph/0504076
7. R. Hagedorn, *CERN preprint* Ref.TH.3684-CERN (1983)
8. M. Biyajima, M. Kaneyama, T. Mizoguchi and G. Wilk, *Euro Phys. J.*, **C40**, 243 (2005)